

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

massing of birds in the south the whole theory would totter. In this connection it is urged that all observers make an unusual effort to make records of abundance during the remaining wintry months as well as during the period of migration, with a view to comparing the numbers of individuals with those which form an average for your vicinity in normal years. The opportunity for learning something more definite about the variations of abundance in birds and its relation to associated phenomena is an unusual one. The editor would be glad to correlate notes which may be sent to him for publication in the June Bulletin.

THE BIRDS OF OHIO: CORRECTION AND NOTE.

LYNDS JONES.

In his Birds of North and Middle America, Prof. Robert Ridgway leaves open the question of the forms of two birds in the Lower Lakes region. He had not seen specimens from the region. One of these is the Long-billed Marsh Wren and the other Traill's Flycatcher. These were entered in my Catalogue of the Birds of Ohio as Telmatodytes palustris and Empidonax traillii respectively, with the note that E. t. alnorum might be looked for as a migrant across the state. Numbers of specimens of each of these forms have since been collected in various parts of the state, and it is now possible to speak with certainty concerning the forms represented in Specimens of the flycatcher have been compared with typical E. traillii from the western states, and have also been submitted to Messrs. Brewster and Oberholser for comparison with larger series of skins. The unanimous conclusion is that Ohio specimens are E. t. alnorum, Alder Flycatcher, and that no E. traillii occur in the state. Likewise, the wrens have been carefully compared with typical T. palustris and prove to be typical and not the Prairie form, T. p. iliacus.